

Variation of sedimentary structures and grainsize over sandwaves

Ad STOLK

Ministry of Transport, Public Works and Water Management, North Sea
Directorate, P.O.Box 5807, 2280 HV Rijswijk, The Netherlands.

Fax.: (31) 70 3900691

Email: a.stolk@dnz.rws.minvenw.nl

Abstract

Sedimentary structures and grainsizes on sandwaves in the Flemish Bank area are studied in relation to the morphology of the sandwaves. There is a difference between sandwaves in deeper water and sandwaves on top of a sand bank. The trough of the sandwaves in deeper water shows bioturbated structures, indicating a less dynamic environment than the top of the sandwaves, which is characterised by migrating megaripples. The sandwaves on top of the bank do not show bioturbation, but primary structures generated by migrating megaripples. In general the grainsize increases from the troughs towards the crests of the sandwaves.

Introduction

Within the framework of the MAST I Resecused project and the MAST II Starfish project an intensive study is done on the sedimentology of a linear sandbank in the southern North Sea. The study concentrates on the Middelkerke Bank in the Flemish Bank area off the Belgian coast.

On top of the Middelkerke Bank and near the northern end of the bank sandwaves are present. The sedimentological structures and the grainsize of the surficial sediments on these sandwaves are studied in relation to the morphology. The determination of the relation between the morphology and the structure and grainsize is important for the interpretation of sediment transport models based on grainsize variations. It is also necessary for the development of good sampling schemes for future sampling programs.

1 Field data

During cruises with RV Mitra and RV Zirfaea in August and October 1993 boxcores (Reineck) and grabsamples (Van Veen) were taken in the sandwave area.

Boxcores were taken on the top, on the steep slope and in the sandwave trough. Lacquer peels were made of the boxcore samples to study the sedimentary structures. In addition to the study of the sedimentary structures over the sandwaves, a detailed grabsampling is carried out along two profiles in the northern part of the Middelkerke Bank area. Along a 1.5 km long profile perpendicular to the sandwave crests (profile Z1, fig.1) 60 samples were taken on a constant distance of 25 metres. Along a 1.5 km long profile in the direction of the maximum current velocities, perpendicular to the megaripples, 22 samples are taken on a constant distance of 50 metres. The grabsamples were used for grainsize determination.

2 Sedimentary structures

To investigate the relation between morphology and sedimentary structures over a sandwave, detailed boxcore sampling is carried out on two sandwaves (fig.1). One sandwave (A) is situated in an area with large sandwaves adjacent to the northern end of the Middelkerke Bank at a water depth of 18 metres (MLLWS). The other sandwave (B) is situated at the top of the bank at a water depth of 7.5 metres (MLLWS). The sandwave in the north (fig.2) shows primary structures due to migrating megaripples on the top. The megaripples on the gentle slope are well developed and reach heights of 90 cm. In the trough of the sandwave the sediments are highly bioturbated. The upper layer on the steep slope shows both primary and bioturbated facies. The sand on top of the sandwave is coarser than in the trough. The height difference between the top and the trough is 5 metres. This difference seems enough to have a dynamic local environment at the top and a less dynamic environment in the trough.

Another reason for the difference can be that the sandwave is situated around a depth of 16 m (MLLWS). This depth is considered to be the boundary between the dynamic and less dynamic environment on the Middelkerke Bank (Stolk, 1993). The 1.3 m high large sandwave on the top of the bank (fig.3) does not show these differences in structure between top and trough. On both places primary structures are present, indicating that the whole sandwave is part of a highly dynamic environment.




Fig.1 Location of studied sandwaves and grabsample profiles

profile Z2




Fig.4 Morphology and grainsize of profile Z2. Grainsize is determined by settling tube (D50 st) and by laser diffraction (D50 ld)

In figure 4 both the morphology and the grainsize of the sand fraction along profile Z2 are shown. Going from SW to NE there is a clear increase in grainsize from the troughs towards the crests of the sandwaves. The same relation is found in the western part of profile Z1 (fig.5). The grain sizes are highly influenced by shell fragments. Remarkable is that on some sandwaves the largest grain sizes are found just below the top at the gentle slope and the smallest values at the top or just below the top at the steep slope. In general this difference in grain size over sandwaves is also found on the Oostdijk Bank (Houthuys, 1990) and outside the bank area in the southern North Sea (Terwindt, 1971; Cameron et al., 1989). In the western part of profile Z1 the morphology is determined by a complex of sandwaves in the swale. The morphology of the eastern part is determined by the northern end of the Middelkerke Bank, where sandwaves are superpositioned on the bank. This morphological difference is also expressed in the grain size. The median grain size in the sandwave field is around 515 (m). On the bank it drops to 420 (m).

profile Z1




Fig.5 Morphology and grainsize of profile Z1. Grainsize is determined by settling tube (D50 st) and by laser diffraction (D50 ld)

Within the area of sandwaves (profile Z2 and the western part of profile Z1) there is a slight trend towards a finer grainsize in deeper water, but there is no significant correlation. Also no correlation is found between grainsize and sandwave height.

Conclusion

The sedimentary structures and the grainsize on sandwaves are related to the morphology. However, the location of the sandwave itself is also an important factor. In deeper water, near the northern end of the bank, the sandwaves show primary, megaripple related structures on the top and bioturbated structures in the trough. This indicates a difference in a dynamic environment on the top and a less dynamic environment in the trough. It is not clear if this difference is a pure local sandwave feature or related to a more general feature in the sand bank area. Sandwaves on top of the sand bank show only primary structures, indicating a highly dynamic environment both on the top and in the trough of the sandwave. In general the grainsize increases from the troughs towards the crests of the sandwaves. Clear correlation between grainsize and water depth or sandwave height are not found.

Reference list

- Cameron, T.D.J., Schüttenhelm, R.T.E. and Laban, C. (1989) Middle and Upper Pleistocene and Holocene stratigraphy in the southern North Sea between 52 and 54 N, 2 to 4 E. In: Henriët, J.P. and De Moor, G. (eds.) *The Quaternary and Tertiary Geology of the Southern Bight, North Sea*. Ministry of Economic Affairs Belgian Geological Survey, 119-135.
- Houthuys, R. (1990) *Vergelijkende studie van de afzettingstructuur van getijdzanden uit het Eoceen en van de huidige Vlaamse Banken*. Aardkundige Mededelingen 5, Leuven University Press, Leuven, 127 p.
- Stolk, A. (1993) Small scale sedimentary structure analysis by boxcoring. In: De Moor, G. and Lanckneus, J. (eds.) *Sediment mobility and morphodynamics of the Middelkerke Bank*. Univ. Gent, CEC, Brussels, 119-132.
- Terwindt, J.H.J. (1971) Sand waves in the Southern Bight of the North Sea. *Marine Geology* 10, 51-67.